

Combination of Dandy — Walker Syndrome and Diabetes Mellitus Type 1

M. V. Novoselova¹, Yu. G. Samojlova¹, N. G. Zhukova¹, O. S. Tonkih²

¹ Siberian State Medical University, Ministry of Healthcare of Russia, Tomsk

² Medical and Diagnostic Center of the International Institute of Biological Systems, Tomsk

Abstract

Frequency of Dandy – Walker syndrome ranges from 3,5 to 12% of malformations of the nervous system, accompanied by hydrocephalus [1]. This syndrome may be asymptomatic and manifests itself in older age, followed by the development of cognitive impairment. In this abstract seems clinical case combination syndrome Dandy – Walker that has been identified when diagnosing the cause of severe cognitive dysfunction in patient with diabetes mellitus type 1.

Key words: Dandy – Walker Syndrome, Diabetes Mellitus Type 1, Cognitive Dysfunction, Quality of Life, Magnetic Resonance Imaging.

References

1. Petrova L. A., Rozanov A. V., Barashnev Ju. I., Panov V. O. Dandy – Walker syndrome in newborns. *Ros. Vestn. Perinatol. Pediat.* 2010. V. 1. P. 25–29 (in Russian).
2. Spennato P., Mirone G., Nastro A. Buonocore M.C., Ruggiero C., Trischitta V., Aliberti F., Cinalli G. Hydrocephalus in Dandy – Walker malformatioSpenn. et al. *Show all Childs Nerv. Syst.* 2011. V. 27. N. 10. P. 1665–1681.
3. Warwick C. T., Reyes B.J., Ayoob M. R., Subit M. Adult diagnosed Dandy –Walker malformation presenting as an acute brainstem event. A case report and review of the literature. et al. *Show W. V. Med. J.* 2008. V. 104. N. 1. P. 25–37.
4. Zhaoyu Gan, Feici Diao, Zili Han, Kanglai Li, Liarong Zheng, Nianhong Guan, Zhuang Kang, Xiaoli Wu, Qinling Wei, Minfeng Cheng, Ming Zhang, Jinbei Zhang. Psychosis and Dandy – Walker complex: report of four cases // et al. *Show General Hospital Psych.* 2012. V. 34. N. 1. P. 102. e7–11.
5. Yamasaki M., Nonaka M., Bamba Y. Yamasaki M., Nonaka M., Bamba Y., Teramoto C., Ban C., Pooh R. K. Diagnosis, treatment, and long-term outcomes of fetal hydrocephalus. et al. *ShSeminars in Fetal & Neonatal Med.* 2012. V. 17. P. 330–335.

Authors

Novoselova Marija Vladimirovna, resident of Department Endocrinology and Diabetology, Siberian State Medical University, Ministry of Healthcare of Russia, Tomsk.

Address: 634050, Russia, Tomsk, Moskovsky tract, 2.

Phone number: +7 (913) 820-06-23. E-mail: novoselova.mariia@yandex.ru

Samojlova Julija Gennad'evna, M. D. Med., Professor of Department Endocrinology and Diabetology, Siberian State Medical University, Ministry of Healthcare of Russia, Tomsk.

Address: 634050, Russia, Tomsk, Moskovsky tract, 2.

Phone number: +7 (913) 826-74-24. E-mail: samoilova_y@inbox.ru

Tonkih Ol'ga Sergeevna, Ph.D. Med., Medical and Diagnostic Center of the International Institute of Biological Systems, Tomsk.

Address: 634049, Russia, Tomsk, Rabochaya st., 21.

Phone number: +7 (923) 424-43-05. E-mail: ostonkih@mail.ru

Zhukova Natal'ja Grigor'evna, M. D. Med., Professor of Department Neurology and Neurosurgery, Siberian State Medical University, Ministry of Healthcare of Russia, Tomsk.

Address: 634050, Russia, Tomsk, Moskovsky tract, 2.

Phone number: +7 (913) 824-62-02. E-mail: znatali@yandex.ru