

Functional Magnetic Resonance Imaging of the Lumbar Spine (Review)

A. V. Bazhin, E. A. Egorova

Moscow State University of Medicine and Dentistry named after A. I. Evdokimov, Ministry of Healthcare of Russia, Department of Radiology

Abstract

The high prevalence of the lumbar spine degenerative changes, significant disability of patients with chronic pain syndrome along with low correlation of radiological and clinical presentation requires searching and implementation of new survey techniques to the diagnostic algorithm. Functional magnetic-resonance imaging provides a new perspective to study the physiology and pathophysiology of degenerative processes. These techniques allow evaluating almost all the structures of the vertebral-motion segments and its positional relationship with anatomical structures of the vertebral canal and intervertebral foramen. In addition, functional magnetic resonance methods permit to detect the signs of instability of soft tissue structures of the spine and neural structures compression during the axial load researches.

Key words: Weight-Bearing Magnetic Resonance Imaging, Axial Load Magnetic Resonance Imaging, Lumbar Spine, Spine Degenerative Disease.

References

1. *Bazhin A. V., Egorova E. A., Kozlov A. E.* The use of weight-bearing Magnetic Resonance Imaging in Lumbar Spine Examination. *Radiology – practice*. 2014. No. 2. P. 6–14.
2. *Bazhin A. V., Egorova E. A., Novoselova E. V., Vasiliev Yu. A., Pakhomov G. A., Berkutov E. S., Durov A. N.* Methods of magnetic resonance imaging with axial load for assessment of lumbar spine under normal and degenerative changes. *MIA Medical Bulletin*. 2015. V. 74. No. 1. P. 49–54.
3. *Shchedrenok V. V., Yakovenko I. V., Anikeev N. V., Sebelev K. I., Moguchaya O. V.* Minimally invasive surgery of degenerative diseases of the spine. St. Petersburg, 2011. 435 p.
4. *Ayturk U. M., Garcia J. J., Puttlitz C. M.* The micromechanical role of the annulus fibrosus components under physiological loading of the lumbar spine. *Journal of Biomechanical Engineering*. 2010. V. 132. No. 6. P. 061007.
5. *Hansen B., Bendix T., Grindsted J., Bliddal H., Hansen P., Christensen R., Riis R., Boesen M.* The effect of low back pain on positional changes in the lumbar lordosis: a cross-sectional comparison with healthy controls. *Insights Imaging*. 2015. No. 6. P. 368
6. *Hebelka H., Brisby H., Hansson T.* Comparison between pain at discography and morphological disc changes at axial loaded MRI in patients with low back pain. *European Spine Journal*. 2014. V. 23. No. 10. P. 2075–2082.
7. *Herkowitz H. N., Garfin S. R., Eismont F. J., Bell G. R., Balderston R. A.* Rothman-Simeone. *The Spine* (6th ed.). Elsevier, 2011. 2000 p.
8. *Kapural L., Kim Ph., Deer T.* Diagnosis, Management, and Treatment of Discogenic Pain. Elsevier, 2011. P. 29–47.

9. Knott P., Mardjetko S., Kim R., Cotter T., Techy F., Rollet M. Comparing axial loaded MRI to standing radiographs in the evaluation of AIS. *Scoliosis*. 2010. No. 5. P. O12.
 10. Lee S. H., Daffner S. D., Wang J. C. Does lumbar disk degeneration increase segmental mobility in vivo? Segmental motion analysis of the whole lumbar spine using kinetic MRI. *J. Spinal. Disord. Tech.* 2014. V. 27. No. 2. P. 111–116.
 11. Martins N., Nunes R. G., Ferreira H., Figueiredo J. P., Barroso L., Mineiro J. D. Impact of patient positioning during MRI of the Lumbar Spine in patients with stenosis: changes in lordosis and canal dimensions. *Insights Imaging*. 2015. No. 6. P. 367–368.
 12. Niggemann P., Kuchta J., Beyer H. K. et al. Spondylolysis and spondylolisthesis: prevalence of different forms of instability and clinical implications. *Spine*. 2011. V. 36. No. 22. P. 1463–1468.
 13. Ottanello C., Giardino A., Giuliani P., Messina C., Di Leo G., Sardanelli F. Differences between supine and orthostatic positions in facet joint (FJ) fluid and spondylolisthesis at lumbar-spine MRI. *Insights Imaging*. 2015. No. 6. P. 368.
 14. Segebarth P. B., Kurd M. F., Haug P. H., Davis R. Routine Upright Imaging for Evaluating Degenerative Lumbar Stenosis: Incidence of Degenerative Spondylolisthesis Missed on Supine MRI. *J. Spinal. Disord. Tech.* 2014. doi: 10.1097/BSD.0000000000000205.
 15. Shymon S. J., Yaszay B., Dwek J. R., Proudfoot J. A., Donohue M., Hargens A. R. Altered disc compression in children with idiopathic low back pain: an upright magnetic resonance imaging backpack study. *Spine*. 2014. V. 39. No. 1. P. 243–248.
 16. Yoder J. H., Peloquin J. M., Song G., Tustison N. J., Moon S. M., Wright A. C., Vresilovic E. J., Gee J. C., Elliott D. M. Internal three-dimensional strains in human intervertebral discs under axial compression quantified noninvasively by magnetic resonance imaging and image registration. *J. Biomech. Eng.* 2014. V. 136. No. 11. doi: 10.1115/1.4028250.
-

Authors

Bazhin Alexander Vladimirovich, Postgraduate of Department of Radiology, Moscow State Medical University of Medicine and Dentistry named after A. I. Evdokimov, Ministry of Healthcare of Russia.
Address: Vucheticha ul., 9a, Moscow, 127206, Russia.
Phone number: +7 (495) 611-01-77. E-mail: avbazhin@yandex.ru

Egorova Elena Alekseevna, M. D. Med., Professor of Department of Radiology, Moscow State Medical University of Medicine and Dentistry named after A. I. Evdokimov, Ministry of Healthcare of Russia.
Address: Vucheticha ul., 9a, Moscow, 127206, Russia.
Phone number: +7 (495) 611-01-77. E-mail: tylsit@mail.ru